

Inferring model parameters in markets with collars

Robert W. Chen¹, Burton Rosenberg¹, & Yi-Tsung Lee²

¹*University of Miami*

²*National Chung Cheng University*

Abstract

Security prices are set by a continuous auction, the rules of which are set by the exchange or by the government. For many exchanges, there is a general free-flow of price information resulting in stock prices which can be modelled by a random walk following a Weiner-Levy process. However, many markets have collars, under which the rules of the auction will not let prices move too rapidly. In this paper we present methods for estimating the volatility of the underlying price data when the true price information is obscured by such collars. Numerical simulations are presented which demonstrate and contrast the methods.

Key words and phrases. Estimation of volatility, market models, market collars.

1 Introduction

Security prices are set by a continuous auction. The rules of the auction are set by the exchange or by the government. For many exchanges, there is a general free-flow of price information resulting in stock prices which can be modelled by a random walk following a Weiner-Levy process, perhaps with an added component of drift depending on the season, investor psychology, or other poorly understood factors. This model of stock prices has been very important both theoretically and practically to the development of option pricing.

However, many markets have collars, under which the rules of the auction will

not let prices move too rapidly. We consider the case where the previous day's closing price is set as the center of a percentage band outside of which the stock will not trade for that day. Since this affects the volatility of stock prices, and options are priced as a function of volatility, the question arises as to the proper pricing of options in markets with collars.

In this paper we present three methods by which the volatility of the ideal price can be estimated by looking only at the exchange price: a method based on the measure of waiting times; a method based on the estimate of the likelihood of an at-market day being followed by an at-limit day (definitions follow); the renewal time method introduced by Chiang and Wei. [2]. In the full paper we provide a rigorous proof of strong consistency for their method.

In addition, we compare the results of these methods to the volatility of the actual stock price and perform simulations on which of the two values are more appropriate for option pricing.

2 Methods

2.1 Definitions

Our ideal stock trace is a sequence of closing prices S_0, S_1, \dots . The inter-day increments are independent, identically distributed random variables with log-normal distribution of common mean μ and variance σ^2 ,

$$Z_i = \ln(S_i/S_{i-1}) \sim N(\mu, \sigma^2).$$

If r is the risk-free rate, then $\mu = r - \sigma^2/2$.

Collars are applied to the ideal price trace S_i to create an observable sequence S_i^* of exchange prices. Given lower and upper collars,

$$\kappa_l < 1 < \kappa_u,$$

we define S_i^* , $i = 0, 1, 2, \dots$ according to,

$$S_i^* = \begin{cases} S_0 & \text{if } i = 0, \\ \kappa_l S_{i-1}^* & \text{if } S_i < \kappa_l S_{i-1}^*, \\ \kappa_u S_{i-1}^* & \text{if } S_i > \kappa_u S_{i-1}^*, \\ S_i & \text{otherwise.} \end{cases}$$

We classify each observation S_i^* as either *at-market* or *at-limit*. Since we cannot always observe S_i , an observation S_i^* will be considered at-market only if it is




Figure 1: Ideal, exchange prices and delta

strictly within that observation's collar. Else we must conclude that the observation is at-limit. Formally, the set of market observations is,

$$T = \{i > 0 \mid (\kappa_l S_{i-1}^*) < S_i^* < (\kappa_u S_{i-1}^*)\} \cup \{0\}$$

and the set of limit observations is the complement,

$$L = T^c = \{i > 0 \mid S_i^* = (\kappa_l S_{i-1}^*) \text{ or } S_i^* = (\kappa_u S_{i-1}^*)\}.$$

We enumerate the indices of T and L in ascending order as T_0, T_1, \dots and L_0, L_1, \dots . Note that by construction $T_0 = 0$, and that L might be empty.

An example trace of 90 day data is given in figure 1. The ideal price, exchange price and their difference are plot, with the difference trace shifted and scaled for display. The simulated stock has zero mean, $\sigma = 47\%$, and the collars were set to $(5/4)\sigma$.

2.2 Estimation of the mean

We estimate the mean μ of S_i and prove some additional lemmas.

If we consider any two consecutive at-market observations, we have a random variable,

$$X_i = \ln(S_{T_i}/S_{T_{i-1}}) = \sum_{j=T_{i-1}+1}^{T_i} Z_j.$$

Since Z_i and Z_j are independent for $i \neq j$, the X_i are independent identically distributed random variables. This is also true that the random variable $T'_i = T_i - T_{i-1}$ are independent and identically distributed.

Define $B_k(a)$ to be the event that the ideal price is above the upper limit of the exchange price for at least k consecutive days given that at day zero the ideal price is a above the zero-th day upper limit,

$$B_k(a) = \{ (a + Z_1 + \dots + Z_j) > j \ln \kappa_u, \text{ for } j = 1, \dots, k \},$$

Lemma 1 Let $\lambda = (\ln \kappa_u - \mu)/\sigma$. If $\lambda > 0$ then for large enough k and $a \geq 0$,

$$\text{Prob}(B_k(a)) < \frac{e^{-k\lambda^2/4}}{\lambda\sqrt{\pi k}}.$$

PROOF: To remain above the exchange limit for k steps requires that the current ideal price exceeds by $k \ln \kappa_u$ the exchange price k days earlier. Since,

$$Y = a + Z_1 + \dots + Z_k \sim N(a + k\mu, k\sigma^2),$$

then,

$$\begin{aligned} \text{Prob}(Y > k \ln \kappa_u) &= \frac{1}{\sqrt{2\pi k\sigma^2}} \int_{k \ln \kappa_u}^{\infty} e^{-(x-k\mu-a)^2/(2k\sigma^2)} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{(\lambda\sqrt{k})-a/(\sigma\sqrt{k})}^{\infty} e^{-x^2/2} dx. \end{aligned}$$

Because $\lambda > 0$, we can select a K such that for all $k \geq K$,

$$(\lambda\sqrt{k}) - a/(\sigma\sqrt{k}) > \lambda\sqrt{k/2} > 0$$

and we can use the approximation for the tail of the normal distribution,

$$\frac{1}{\sqrt{2\pi}} \int_x^{\infty} e^{-y^2/2} dy < \frac{1}{x\sqrt{2\pi}} e^{-x^2/2}$$

for $x > 0$, see Feller, Vol. 1, Lemma VII.1.7. Since the bound is increasing with decreasing x , for simplicity of notation we set $x = \lambda\sqrt{k/2}$ to complete the proof.

Lemma 2 If $\ln \kappa_l < \mu < \ln \kappa_u$ then $E(T_1)$ and $E(T_1^2)$ are bound.

PROOF: A sequence of days which stay outside of the market will consist of segments of days in one collar alternating with segments of days in the other collar.

Consider now the random variable B^u which gives the number of days we can stay within the upper collar. Note that $B_k^u = \bigcap_{j=1}^k B_j$. Using the previous lemma,

$$\begin{aligned} E(B^u) &< \sum_{k=0}^{\infty} \text{Prob}(B_k) \\ &< C_N + \sum_{k=N}^{\infty} \frac{e^{-k\lambda^2/4}}{\lambda\sqrt{\pi k}} < \infty, \end{aligned}$$

where $\lambda = (\ln \kappa_u - \mu)/\sigma$. An similar bound exists for expectation time in the lower collar $E(B^l)$.

Now consider the random variable C which gives the number of passages from upper to lower collars before the next consecutive day at-market. The probability to pass from one collar to the other must be at least,

$$p_c = \text{Prob}(Z_j \geq (\ln \kappa_u - \ln \kappa_l)).$$

Considering the various cases by which a random walk might stay outside of either collar, we arrive at, essentially, a sum over i changes from upper to lower of walks of approximate length weighted by the probability of i changes of collars,

$$\begin{aligned} E(T_1) &\approx \text{Prob}(\ln \kappa_l < Z_1 < \ln \kappa_u) + \sum_{i=1}^{\infty} (p_c^{i-1})(i/2)(E(B^u) + E(B^l) + i) \\ &< K_c + K \sum_{i=1}^{\infty} i^2 p_c^i < \infty. \end{aligned}$$

The proof for $E(T_1^2)$ is similar.

Lemma 3 $E(X_{T_1}) = \mu E(T_1)$.

PROOF: It is now more convenient to work with increments $\mu + \sigma Z_i$ with respect to normalized random variables Z_i .

$$\begin{aligned} E(X_{T_1}) &= E\left(\sum_{j=1}^{T_1} \mu + \sigma Z_j\right) \\ &= \sum_{k \geq 1} \int_{\{T_1=k\}} \sum_{j=1}^k (\mu + \sigma Z_j) dP \\ &= \sum_{k \geq 1} k \mu P(T_1 = k) + \sigma \sum_{k \geq 1} \sum_{j=1}^k \int_{\{T_1=k\}} Z_j dP \end{aligned}$$

The first term on the right-hand side is obviously $\mu E(T_1)$. Rearranging the sum of the second term on the right hand side,

$$\begin{aligned} \sigma \sum_{k \geq 1} \sum_{j=1}^k \int_{\{T_1=k\}} Z_j dP &= \sigma \sum_{j \geq 1} \sum_{k \geq j} \int_{\{T_1=k\}} Z_j dP \\ &= \sigma \sum_{j \geq 1} \int_{\{T_1 \geq j\}} Z_j dP \\ &= \sigma \sum_{j \geq 1} \left(E(Z_j) - \int_{\{T_1 < j\}} Z_j dP \right) \end{aligned}$$

Since Z_j and $\{T_1 < j\}$ are independent,

$$E(Z_j) - \int_{\{T_1 < j\}} Z_j dP = E(Z_j)(1 - P(T_1 < j)) = 0$$

because $E(Z_j) = 0$. We conclude that the second term is zero.

Since $E(X_{T_1})$ and $E(T_1)$ are finite, the Strong Law of Large Numbers gives us strong consistency for the sample expectations of X_{T_1} and T_1 ,

$$\begin{aligned} E(\widehat{X}_{T_1}) &= (1/n) \sum_{i=1}^n X_i \\ E(\widehat{T}_1) &= (1/n) \sum_{i=1}^n (T_i - T_{i-1}) = T_n/n \end{aligned}$$

and a strongly consistent approximation $\hat{\mu}$ to μ ,

$$\hat{\mu} = E(X_{T_1})/E(T_1) = (\ln S_{T_n} - \ln S_0)/T_n$$

where $n = |T|$ is the number of days observed at-market in our sample.

Theorem 1 *Under the given assumptions $\hat{\mu}$ is strongly consistent.*

2.3 Estimation of σ : the waiting time method

To estimate the variance, we look at the waiting time for the next at-limit observation beginning from an at-market observation. Define T' to be the first at-market

day in a sequence of at market days, and L' to be the first at-limit day in a sequence of at-limit days,

$$T' = \{i \in T \mid i > 1, i - 1 \notin T\} \cup \{0\}$$

$$L' = \{i \in L \mid i - 1 \in T\}$$

Enumerate these indices as T'_1, T'_2, \dots, T'_m and L'_1, L'_2, \dots, L'_m . Pairing them up we calculate the i -th waiting time $W_i = L'_i - T'_i$. The W_i are independent identically distributed random variables with geometric distribution $\{pq^{k-1}\}$ where q is the probability that Z_i falls within the collar $\ln \kappa_l < Z_i < \ln \kappa_u$. Hence,

$$E(W_1) = 1/(1 - q) < \infty.$$

This gives us the implicit relation between $E(W_1)$ and σ ,

$$1 - \frac{1}{E(W_1)} = \frac{1}{\sqrt{2\pi}} \int_{(\ln \kappa_l - \mu)/\sigma}^{(\ln \kappa_u - \mu)/\sigma} e^{-x^2/2} dx.$$

Since $E(W_1) < \infty$, the Strong Law of Large Numbers allows that the sample mean $\widehat{E(W_1)}$ is a strongly consistent estimate for $E(W_1)$,

$$\widehat{E(W_1)} = (1/m) \sum_{i=1}^m W_i.$$

We use numerical methods to solve for $\hat{\sigma}$ given $\widehat{E(W_1)}$. Due to the continuity of this relation, that $\widehat{E(W_1)}$ is a strongly consistent estimate for $E(W_1)$ implies that $\hat{\sigma}$ is a strongly consistent estimate for σ .

Theorem 2 *Under the given assumptions $\hat{\sigma}$ is strongly consistent.*

2.4 Estimation of σ : the transition counting method

As referred to in the previous section, we can view the outcome of what follows an at-market day as the flip of a biased coin. The bias on this coin relates to the area under the gaussian distribution which might put the next market step into the limit region — and this is related monotonically to the variance. By observing the ratio of at-market days followed by at-market days versus by at-limit days, this bias can be estimated.

Define R to be the set of indices such that two consecutive days are at market and S to be the set of indices such that the first day is at market and the next is at limit,

$$T_{MM} = \{i \mid (i \in T) \wedge (i + 1 \in T)\},$$

$$T_{ML} = \{i \mid (i \in T) \wedge (i + 1 \in L)\}.$$

We are flipping a coin which comes up “limit” with probability $|T_{ML}|/(|T_{ML}| + |T_{MM}|)$. Therefore,

$$E(W) = 1 + (|T_{MM}|/|T_{ML}|).$$

This formula agrees with the formula for $E(W)$ of the waiting time method if the sequence of observations ends on an at-limit day. Else the estimate of $E(W)$ will be slightly lower, as the sequence of ending observations will bias downward the estimation of the probability of hitting the limit.

Because this method agrees so closely with the method of waiting times, we will not consider it further.

2.5 Estimation of σ : the renewal time method

Recall that our at-market days have indices T_0, T_1, \dots . The renewal time from one at-market day to another is $T'_i = T_{i-1} - T_i$, for $i = 1, 2, \dots$. The T'_i are i.i.d. random variables. The change in log market price during interval T'_i is,

$$X_{T'_i} = \sum_{j=T_{i-1}-1}^{T_i} (\mu + \sigma Z_j)$$

where the Z_i are i.i.d. standard gaussian normal random variables.

Lemma 4 $E((X_{T_1} - \mu T_1)^2) = \sigma^2 E(T_1)$.

PROOF: Proof in full paper.

Theorem 3 *The estimate,*

$$\hat{\sigma}^2 = \frac{\sum_{i=1}^n (X_{T'_i} - \hat{\mu} T'_i)^2}{\sum_{i=1}^n T'_i}$$

is strongly consistent.

PROOF: Proof in full paper.

3 Results

In graphs in figures 2 through 4 plot results of the estimation of σ for the waiting time method, the renewal time method, and the direct calculation on the observed price stream, with this data displayed as box charts, left to right as listed.

For each trial, a random walk of 250 days was generated, with $\mu = 0$ and $\sigma = 1$. The upper and lower collars equal a common κ , where these collars are applied to the log data,

$$\ln \kappa_l = \ln \kappa_u = \kappa.$$

Various ratios κ/σ are tried. A plot of 100 trials with a bisected box indicating the median and one quartile above and below the median is given.

For lower κ/σ , the waiting time method shows greater accuracy (median is closer to 1) and less dispersion (quartiles are closer to median). By $\kappa/\sigma = 2$ the two methods perform about equally, and above that the renewal time method tends to excel. In this region, hitting a collar is rare, so we do not get enough waiting time events to accurately measure σ .

References

- [1] Robert Cast, André Lapiéd, *Fondements Microéconomique de la Théorie des Marchés Financiers*, Economica, Paris, 1992
- [2] R. Chiang, K. Wei, *GMM and MLE approaches for estimation of volatility and regression models when daily prices are subject to price limits*, manuscript.
- [3] William Feller, *An Introduction to Probability Theory and its Applications*, Volume 2, John Wiley and Sons, 1996.
- [4] John C. Hall, *Options, Futures, and Other Derivatives*, Prentice-Hall, 1997.




Figure 2: $\kappa/\sigma = 1/2$




Figure 3: $\kappa/\sigma = 1$




Figure 4: $\kappa/\sigma = 2$